

* CONSTITUTIONAL CRITERIA									
FEVER	<p>ONE of the following criteria MUST be met:</p> <ol style="list-style-type: none"> 1. Single temperature more than 37.8°C 2. Repeated oral temperatures of more than 37.2°C or rectal temperatures more than 37.5°C 3. Single temperature more than 1.1°C increase over baseline of non-illness temperature collected from any site <p><i>There is insufficient evidence to indicate a specific time frame for evaluating repeated temperatures using fever criterion 2. It is suggested repeated temperatures be collected within no more than 48 hours of each other</i></p>								
LEUKOCYTOSIS	>11 x 10 ⁹ leukocytes/L								
ACUTE CHANGE in MENTAL STATUS from BASELINE USING the CONFUSION ASSESSMENT METHOD	<p>ALL FOUR criteria must be present:</p> <ol style="list-style-type: none"> 1. Evidence of acute change in mental status 2. Fluctuating course: behaviour fluctuating (e.g., coming and going, or changing in severity during the assessment) 3. Inattention: difficulty focusing attention (e.g., unable to keep track of discussion or easily distracted) 4. EITHER A OR B: <ol style="list-style-type: none"> A. Disorganized thinking B. Altered level of consciousness: level of consciousness is described as different from baseline (e.g., hyper-alert, sleepy, drowsy, difficult to arouse, non- responsive) 								
ACUTE FUNCTIONAL DECLINE	<p>A new three-point increase in total Activities of Daily Living (ADL) score (range 0-28) from baseline, based on the following seven ADL items, each scored from 0 (independent) to 4 (total dependence)</p> <table style="width: 100%; border: none;"> <tr> <td style="width: 50%;">1. Bed mobility</td> <td style="width: 50%;">2. Transfer</td> </tr> <tr> <td>3. Dressing</td> <td>4. Locomotion within LTC home</td> </tr> <tr> <td>5. Toilet use</td> <td>6. Personal hygiene</td> </tr> <tr> <td>7. Eating</td> <td></td> </tr> </table> <p><i>Note: This information may not be readily available. If unable to appropriately determine rely on the other indicators above to meet the constitutional criteria.</i></p>	1. Bed mobility	2. Transfer	3. Dressing	4. Locomotion within LTC home	5. Toilet use	6. Personal hygiene	7. Eating	
1. Bed mobility	2. Transfer								
3. Dressing	4. Locomotion within LTC home								
5. Toilet use	6. Personal hygiene								
7. Eating									

INSTRUCTIONS for USE

- | | |
|--|---|
| <ol style="list-style-type: none"> 1 Add all residents with respiratory symptoms to this line list. 2 No white out to be used on line list. 3 If error made, strike out and use a new line. 4 Exclude cases with existing or chronic respiratory conditions who are at baseline. 5 Start a new line list sheet for each new month. 6 Retain hard copy for record retention. 7 Ensure notification to ICSA/ICP continues by facility identified process. | <ol style="list-style-type: none"> 8 For questions or concerns, contact facility ICSA. 9 Print on double sided legal paper requesting the printer to flip pages on the short side. 10 First nurse to make assessment, put initials at end of row; Additional symptoms can be added on the same row later using new nurses' initials in the appropriate column. |
|--|---|