
















Microorganism, Infectious Disease Table Highlights

MICROORGANISM INFECTIOUS DISEASE	TYPE OF PRECAUTIONS	SPECIAL CONSIDERATIONS	APPROPRIATE ROOM	ROOM
Antimicrobial Resistant Gram Negative Bacteria	Contact 	Implement Contact Precautions for patients positive for or flagged as AMR GNB.	Single room	
Bed Bugs (cimex lectularius)	Routine Practices Contact if more than 1 bed bug observed	Bag personal belongings. Remove privacy curtains. Contact Housekeeping/Facility Management/ Environmental Services for management.	Single room	
Carbapenemase-Producing Enterobacteriaceae positive (CPE POS) Or CPE suspect (CPE SUS)	Containment 	Do not collect specimen if positive. Suspect: Collect specimens Notify site ICP; leave a message after hours. Consult site ICP BEFORE transport or cohorting if single rooms are limited. Or contact: Dr. Lo @ 204-932-6538 at SBH; Dr. Embil @ 204-931-9538 at other sites; Attending Ped. ID MD: 204-787-2071: pediatric cases.	Single room	
Clostridioides difficile Infection (<i>C. difficile</i> /CDI)	Contact 	Perform hand hygiene at the point of care using soap and water. If a sink is not available at the point-of-care, use alcohol-based hand rub. Perform hand hygiene with soap and water as soon as a sink is available.	Single room	
Chickenpox	Airborne/Contact 	Patient wears a procedure mask if not in appropriate room. HCWs, roommates and caregivers should be immune to chickenpox. People with known immunity do not need an N95 respirator to enter the room.	Airborne Infection Isolation Room (AIIR) or single, door closed	
ILI (influenza like illness)	Droplet/Contact 	Patient wears a procedure mask if not in appropriate room. If private room is unavailable, consider cohorting patients during outbreaks . Do not cohort with high-risk roommates.	Single room at least 2 metres from other patients	
Immunocompromised Patient (Severe)	Protective Precautions 	Have had an allogeneic hematopoietic stem cell transplant OR Absolute Neutrophil Count (ANC) is expected to be below 0.5 x 10E9/L for 7 days or more	Positive pressure OR Single room, door closed	
Immunocompromised Patient (Moderate to Severe)	Modified Protective Precautions 	Have NOT had an allogeneic hematopoietic stem cell transplant AND Absolute Neutrophil Count (ANC) is NOT expected to be below 0.5 x 10E9/L for 7 days	Single room, door open	
Lice	Routine Practices	Routine Practices plus gloves for direct patient contact only	Any	

Microorganism, Infectious Disease Table Highlights

Measles (Rubeola)	Airborne		Patient wears a procedure mask if not in appropriate room. Reportable Disease: Reported by diagnosing HCW. See Reporting of a Communicable Disease to MB Health by IP&C in Hospitals Operational Directives . Contact Site ICP.	Airborne Infection Isolation Room (AIIR) or single, door closed
Meningitis (unknown type)	Adults & > 5 years Droplet Under 5 Droplet/Contact		When type of meningitis (virus or <i>Neisseria meningitidis</i> or other bacteria), is identified refer to Microorganism, Infectious Disease Table or Meningitis Protocol for more information.	Single room at least 2 meters from other patients
Meningitis (Viral)	Adults: Routine Practices		Contact Precautions : children under 5 years, incontinent person whose feces cannot be contained; People who contaminate the environment	Adult: any room Child under 5: Single room
MRSA SUS (Suspect)	Routine Practices		Collect specimens as for MRSA POS (positive)	Any
MRSA POS (Positive)	Contact		Collect specimens from: nares AND open wounds/lesions/incisions/invasive device insertion sites	Single room
Mumps (positive or exposed and susceptible)	Droplet		Immune people do not need PPE. HCWs, roommates and visitors should be immune to mumps. Non-immune HCWs should not enter room if immune caregivers are available. Contact Site ICP.	Single room at least 2 metres from other patients
Necrotizing Fasciitis	Contact		Duration of precautions: Until 24 hours of appropriate antimicrobial therapy received	Single room
Scabies	Contact		Wash clothes and bedding in hot water, dry clean or seal in a plastic bag, and store for 1 week. Household contacts and exposed staff should be treated.	Single room
Disseminated Shingles (Varicella Zoster)	Airborne/Contact		Patient wears a procedure mask if not in appropriate room. HCWs, roommates and caregivers should be immune to chickenpox. Non-immune HCWs do not enter room if immune caregivers available. If non-immune person enters room, for exceptional circumstances, wear N95 respirator.	Airborne Infection Isolation Room (AIIR) or single, door closed
Tuberculosis	Airborne		Patient wears a procedure mask if not in appropriate room. Reportable Disease: See Reporting of a Communicable Disease to MB Health by IP&C in Hospitals Operational Directives . See Respiratory Protection for Aerosol Generating Medical Procedures (AGMPs) . Contact Site ICP	Airborne Infection Isolation Room (AIIR) or single, door closed