For Asymptomatic/Prophylactic use: For residents with CrCl 10-30 mL/min

Oseltamivir 30 mg po every other day for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl 10-30 mL/min

Oseltamivir 30 mg po every other day for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl 10-30 mL/min

Oseltamivir 30 mg po every other day for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl 10-30 mL/min

Oseltamivir 30 mg po every other day for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl 10-30 mL/min

Oseltamivir 30 mg po every other day for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl 10-30 mL/min

Oseltamivir 30 mg po every other day for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl 10-30 mL/min

Oseltamivir 30 mg po every other day for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl 10-30 mL/min

Oseltamivir 30 mg po every other day for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl 10-30 mL/min

Oseltamivir 30 mg po every other day for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCI 10-30 mL/min

Oseltamivir 30 mg po every other day for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms. For Asymptomatic/Prophylactic use: For residents with CrCI 10-30 mL/min

Oseltamivir 30 mg po every other day for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCI 10-30 mL/min

Oseltamivir 30 mg po every other day for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl 10-30 mL/min

Oseltamivir 30 mg po every other day for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl 10-30 mL/min

Oseltamivir 30 mg po every other day for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl 10-30 mL/min

Oseltamivir 30 mg po every other day for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl 10-30 mL/min

Oseltamivir 30 mg po every other day for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCI 10-30 mL/min

Oseltamivir 30 mg po every other day for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCI 10-30 mL/min

Oseltamivir 30 mg po every other day for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl 10-30 mL/min

Oseltamivir 30 mg po every other day for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCI 10-30 mL/min

Oseltamivir 30 mg po every other day for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms. For Asymptomatic/Prophylactic use: For residents with CrCl 10-30 mL/min

Oseltamivir 30 mg po every other day for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCI 10-30 mL/min

Oseltamivir 30 mg po every other day for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl 10-30 mL/min

Oseltamivir 30 mg po every other day for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl 10-30 mL/min

Oseltamivir 30 mg po every other day for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl 10-30 mL/min

Oseltamivir 30 mg po every other day for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl 10-30 mL/min

Oseltamivir 30 mg po every other day for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl 10-30 mL/min

Oseltamivir 30 mg po every other day for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl 10-30 mL/min

Oseltamivir 30 mg po every other day for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl 10-30 mL/min

Oseltamivir 30 mg po every other day for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl 10-30 mL/min

Oseltamivir 30 mg po every other day for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.