For Asymptomatic/Prophylactic use: For residents with CrCl greater than 30 mL/min up to 60 mL/min

Oseltamivir 30 mg po once daily for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl greater than 30 mL/min up to 60 mL/min

Oseltamivir 30 mg po once daily for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl greater than 30 mL/min up to 60 mL/min

Oseltamivir 30 mg po once daily for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl greater than 30 mL/min up to 60 mL/min

Oseltamivir 30 mg po once daily for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl greater than 30 mL/min up to 60 mL/min

Oseltamivir 30 mg po once daily for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl greater than 30 mL/min up to 60 mL/min

Oseltamivir 30 mg po once daily for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl greater than 30 mL/min up to 60 mL/min

Oseltamivir 30 mg po once daily for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl greater than 30 mL/min up to 60 mL/min

Oseltamivir 30 mg po once daily for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl greater than 30 mL/min up to 60 mL/min

Oseltamivir 30 mg po once daily for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl greater than 30 mL/min up to 60 mL/min

Oseltamivir 30 mg po once daily for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl greater than 30 mL/min up to 60 mL/min

Oseltamivir 30 mg po once daily for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl greater than 30 mL/min up to 60 mL/min

Oseltamivir 30 mg po once daily for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl greater than 30 mL/min up to 60 mL/min

Oseltamivir 30 mg po once daily for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl greater than 30 mL/min up to 60 mL/min

Oseltamivir 30 mg po once daily for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl greater than 30 mL/min up to 60 mL/min

Oseltamivir 30 mg po once daily for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl greater than 30 mL/min up to 60 mL/min

Oseltamivir 30 mg po once daily for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl greater than 30 mL/min up to 60 mL/min

Oseltamivir 30 mg po once daily for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl greater than 30 mL/min up to 60 mL/min

Oseltamivir 30 mg po once daily for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl greater than 30 mL/min up to 60 mL/min

Oseltamivir 30 mg po once daily for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl greater than 30 mL/min up to 60 mL/min

Oseltamivir 30 mg po once daily for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl greater than 30 mL/min up to 60 mL/min

Oseltamivir 30 mg po once daily for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl greater than 30 mL/min up to 60 mL/min

Oseltamivir 30 mg po once daily for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl greater than 30 mL/min up to 60 mL/min

Oseltamivir 30 mg po once daily for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl greater than 30 mL/min up to 60 mL/min

Oseltamivir 30 mg po once daily for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl greater than 30 mL/min up to 60 mL/min

Oseltamivir 30 mg po once daily for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl greater than 30 mL/min up to 60 mL/min

Oseltamivir 30 mg po once daily for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl greater than 30 mL/min up to 60 mL/min

Oseltamivir 30 mg po once daily for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl greater than 30 mL/min up to 60 mL/min

Oseltamivir 30 mg po once daily for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl greater than 30 mL/min up to 60 mL/min

Oseltamivir 30 mg po once daily for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.

For Asymptomatic/Prophylactic use: For residents with CrCl greater than 30 mL/min up to 60 mL/min

Oseltamivir 30 mg po once daily for 10 days or until outbreak declared over, whichever occurs first. Contact prescriber immediately to consider treatment doses if resident develops influenza signs/symptoms.