

ROUTINE PRACTICES PROTOCOL

WINNIPEG REGIONAL HEALTH AUTHORITY
INFECTION PREVENTION AND CONTROL MANUAL

The primary goal of Infection Prevention and Control programs is to reduce the risk of acquiring a healthcare-associated infection (HAI) to a minimum level; zero risk is not possible in every circumstance but should nevertheless be the ultimate goal. The consequences of cross-transmission of microorganisms (germs) must be balanced against the consequences (adverse effects and cost) of precautions taken.

Routine Practices are the foundation for preventing the transmission of microorganisms during care in all healthcare settings. It is a comprehensive set of Infection Prevention and Control (IP&C) measures developed for use in the routine care of **ALL PERSONS** at **ALL TIMES** in **ALL HEALTHCARE SETTINGS (acute, long-term, or community care)**. Routine Practices aim to minimize or prevent healthcare-associated infections in everyone in the healthcare setting including the patient/resident/client (P/R/C), all staff, visitors, Designated Caregivers, contractors, and so on. Following Routine Practices can reduce the transmission of microorganisms in all healthcare settings.

All staff (physicians, nurses, allied health staff, support staff, students, volunteers and others) are responsible for complying with Routine Practices and for tactfully calling infractions to the attention of those whom have missed an opportunity. P/R/Cs and all visitors have a responsibility to follow Routine Practices. Teaching those receiving care and visitors the basic principles of Routine Practices (e.g., hand hygiene, use of personal protective equipment) is the responsibility of all staff. **No one is exempt from following Routine Practices.**

Consistent use of Routine Practices is expected for the care of all persons at all times in all healthcare setting: in hospital, long term, or community care. **It is important to follow Routine Practices at all times for all P/R/C in all healthcare settings** as microorganisms can be transmitted from both symptomatic and asymptomatic people.

TABLE OF CONTENTS

1. POINT OF CARE RISK ASSESSMENT (PCRA)	4
1.1. How to Perform a Point of Care Risk Assessment	4
1.2. Using Control Measures After Performing a PCRA	5
2. HAND HYGIENE	6
2.1. Alcohol-Based Hand Rub (ABHR)	7
2.2. Hand Washing	7
2.3. Care Environments	8
2.4. Indications and Moments for Hand Hygiene During Healthcare Activities	9
3. THE 4 MOMENTS OF HYGIENE	9
3.1. MOMENT 1: BEFORE INITIAL PATIENT/RESIDENT/CLIENT (P/R/C) CONTACT OR P/R/C ENVIRONMENT CONTACT	9
3.2. MOMENT 2: BEFORE ASEPTIC/CLEAN PROCEDURES	10
3.3. MOMENT 3: AFTER BODY FLUID EXPOSURE RISK	10
3.4. MOMENT 4: AFTER PATIENT/RESIDENT/CLIENT CONTACT OR PATIENT/RESIDENT/CLIENT ENVIRONMENT CONTACT	11
3.5. HAND HYGIENE TECHNIQUES	12
3.5.1. Using an Alcohol-Based Hand Rub (ABHR)	12
3.5.2. Using Soap and Water	12
3.5.3. When Hands are Visibly Soiled and Liquid/Gel Soap and Running Water is Not Available	13
3.6. Factors that Reduce Effectiveness of Hand Hygiene	13
3.6.1. Condition of the Hands	13
3.6.2. Nails	14
3.6.3. Nail Polish	14
3.6.4. Artificial Nails or Nail Enhancements	14
3.6.5. Jewelry	15
3.6.6. Upper Extremity Support Devices	15
3.6.7. Other Obstacles to Effective Hand Hygiene	15
3.6.8. Hand Drying (paper towel, air dryers)	16
3.6.9. Lotions and Creams	16
3.6.10. Dispensers	17

4. SOURCE CONTROL	17
4.1. Respiratory Etiquette/Respiratory Hygiene.....	18
4.2. Triage	18
4.2.1. Emergency Rooms and Acute Assessment Settings	18
4.2.2. Ambulatory Care/Clinic Settings	19
4.3. Early Diagnosis and Treatment	19
4.4. Spatial Separation	19
4.5. Aerosol-Generating Medical Procedures (AGMPs)	20
5. ACCOMODATION AND PLACEMENT	23
5.1. Options for P/R/C Placement and Room Sharing.....	23
5.2. Risk Factors for Transmission from the Infection P/R/C?	23
5.3. Priority for Single Rooms Goes to Those:	24
5.4. Factors to be considered with shared rooms include:.....	24
5.5. Cohorting.....	24
5.6. The Use of Airborne Isolation Rooms (AIIRs).....	24
6. FLOW	25
6.1. Flow and Additional Precautions	26
6.2. Ambulatory Care/Clinic Setting.....	26
7. ASEPTIC TECHNIQUE	26
7.1. Recommendations for Injection Safety Include	27
7.2. Aseptic Technique for Invasive Procedures and Handling Injectable Products	27
7.3. Single Dose Vials	28
7.4. Multi-Dose Vials.....	28
7.5. Single P/R/C Multi-Use Devices	29
7.6. Injecting Material and Placing a Catheter into the Spinal Canal or Subdural Space ..	29
7.7. Insertion of Central Venous Catheters.....	29
7.8. Insertion of Peripheral Venous Catheters or Peripheral Arterial Lines.....	29
7.9. Storage, Assembly or Handling Components of Intravenous (IV) Delivery Systems..	30
8. PERSONAL PROTECTIVE EQUIPMENT	30
8.1. Gloves	31
8.1.1. Appropriate Glove Use.....	31
8.2. Long-Sleeved Gowns and Other Apparel.....	32

8.2.1. Appropriate Gown Use.....	33
8.3. Facial Protection.....	34
8.3.1. (Medical) Masks.....	34
8.3.2. Eye Protection.....	35
8.4. Respiratory Protection	36
9. SPECIMEN COLLECTION	36
9.1. Process	37
10. SHARPS SAFETY & PREVENTION OF BLOODBORNE TRANSMISSION	38
11. MANAGEMENT OF THE PATIENT/RESIDENT/CLIENT CARE ENVIRONMENT	39
11.1. Cleaning of Environment	39
11.2. Cleaning and Disinfection of Non-Critical Patient/Resident/Client Care Equipment ...	40
11.2.1. In Home Care Settings.....	41
11.3. Handling of Linen.....	42
11.4. Handling of Waste	43
11.5. Handling of Dishes	43
11.6. Handling of Deceased Bodies	44
12. VISITOR AND DESIGNATED CAREGIVER MANAGEMENT AND EDUCATION	44
13. REFERENCES	46
Appendix A.....	47

1. POINT OF CARE RISK ASSESSMENT (PCRA)

Prior to every interaction with the patient, resident or client (P/R/C); all healthcare staff are responsible to assess the task/care to be performed, the clinical presentation of the P/R/C, physical state of the environment and the healthcare setting, to determine the infectious risk to themselves, those receiving care, Designated Caregivers, visitors, and staff. A Point of Care Risk Assessment (PCRA) is a tool for staff to use before each interaction to determine appropriate action and control measures needed to minimize the risk to staff, the P/R/C and others in the healthcare (external) environment.

**A Point of Care Risk Assessment (PCRA)
should be used by both clinical and non-clinical staff**

1.1. How to Perform a Point of Care Risk Assessment

To perform a PCRA, consider infection transmission risk for the specific:

- Interaction/task
- Environment
- P/R/C, and
- Health care worker

When each staff member performs a [PCRA](#), they must determine the risk of exposure and potential for the spread of microorganisms (germs) during interactions with those receiving care. Examples of factors to consider include:

- ? What kind of contact will I have with the P/R/C (prolonged or frequent direct care)?
- ? What is the health status of the P/R/C? Are they showing signs and symptoms of infection (i.e., coughing, sneezing, respiratory secretions)? Are they immunocompromised?
- ? Is the P/R/C cooperative? Do they understand what is happening?
- ? Will there be a risk of splashes or sprays of blood or body fluids during the task(s) or procedure(s)? Is there a risk of exposure to secretions and excretions, non-intact skin, or mucous membranes?
- ? If the P/R/C has diarrhea, is he/she continent? If incontinent, can stool be contained in a diaper or incontinent product?
- ? Is the P/R/C able and willing to perform hand hygiene and/or wear a medical mask (procedure or surgical mask) if required?
- ? Will an aerosol generating medical procedure (AGMP) be involved?
- ? Is the P/R/C in a shared room/treatment space?

1.2. Using Control Measures After Performing a PCRA


After assessing the status of the P/R/C, the task/procedure, and the care environment, use control measures to lower the chance of spreading potentially harmful microorganisms. Control measures may include:

- Hand hygiene (using alcohol-based hand rub at point of care)
- Placement and accommodation of the P/R/C:
 - Place those with suspected or confirmed airborne infection (e.g., measles or tuberculosis) into an airborne infection isolation room (AIIR) with the door closed
 - Implement strategies to reduce aerosol production in AGMP's (see: [AGMP section 4.5](#))
 - Give priority to those with uncontained wound drainage or uncontained diarrhea into a single room
 - If sharing a room, spatial separation (2m) between beds is ideal and shared equipment must be cleaned between uses.
- Treating an active infection
- Selecting roommates for shared rooms or for transport in shared ambulances (and other types of transportation e.g., air ambulances, taxis), consider the infection risk posed from a P/R/C or posed to the P/R/C
- Consider the immune status of P/R/C who may potentially be exposed to certain infections (e.g., measles, mumps, rubella, varicella)
- Flow (movement) of the P/R/C
 - Restrict movement of symptomatic P/R/C within the specific care area/facility or outside the facility as appropriate for the suspected or confirmed infection/colonization.
- Work assignment: consider the immune status of staff who will potentially be exposed to certain infections (e.g., measles, mumps, rubella, varicella)
- Personal protective equipment (PPE) selection:
 - Use personal protective equipment appropriate to the suspected or confirmed infection/colonization.
- Aseptic technique for invasive procedures such as injections, IV insertions, etc.
- Cleaning and disinfecting non-critical care equipment and the environment.
- Handling of linen and waste
- Restricting visitor access where appropriate
- Assess need for implementing, maintaining or discontinuing Additional Precautions.

2. HAND HYGIENE

Hand hygiene (HH) is a comprehensive term that applies to hand washing, hand antisepsis and to actions taken to maintain healthy hands and fingernails. Hand hygiene is a core element of safe care for the prevention of infections and preventing the spread of microorganisms (germs), including antimicrobial resistant organisms (AROs). Hand hygiene includes cleaning hands with soap and water or alcohol-based hand rub in order to remove microorganisms. HH includes [surgical hand antisepsis](#).^{13.4}

The most common way microorganisms are spread in any healthcare setting is from the hands of health care staff to patient/resident/client (P/R/C), either directly or indirectly. During the delivery of health care, staff constantly touch surfaces and substances including inanimate objects, a P/R/C's intact or non-intact skin, mucous membranes, food, waste, body fluids and the staff member's own body (e.g., hair). All of these can carry potentially harmful microorganisms that can be passed on to someone else if proper HH is not performed. The spread of microorganisms can result in **healthcare-associated infections** (HAIs).



Healthcare workers should teach those receiving care, Designated Caregivers and visitors about HH. The benefits of the general public participating in hand hygiene should not be underestimated. HH education and accessible HH product for the general public is encouraged.

**In healthcare settings,
 hand hygiene is the single most important way to
 prevent infections**

2.1. Alcohol-Based Hand Rub (ABHR)

- Use of alcohol-based hand rub (ABHR) has been shown to reduce healthcare-associated infection rates
- **ABHR is the preferred hand hygiene method** and should be used at point-of-care unless hands are visibly soiled
- ABHR is appropriate to use when caring for P/R/C with *Clostridioides difficile*, except in outbreak or hyperendemic (sustained high rate) settings, then handwashing with soap and water is recommended^{13.11}.
- ABHR is faster and more effective than washing hands (even with an antibacterial soap) when hands are not visibly soiled
- ABHRs:
 - ✓ Provide for a rapid kill of most transient microorganisms
 - ✓ Are not to be used with water
 - ✓ Contain emollients to reduce hand irritation
 - ✓ Take less time than washing with soap and water
- **DO NOT** use ABHR with water as it will reduce its effectiveness
- **DO NOT** use ABHR immediately after hand washing with soap and water as it will result in more hand irritation
- ABHR available for healthcare settings ranges in concentration from 60 to 90% alcohol.

2.2. Hand Washing

Hand washing with soap and running water must be performed when hands are visibly soiled. Antimicrobial soap may be considered for use in critical care settings such as intensive care units and burn units, but is not required and not recommended in other care areas.

- If the P/R/C bathroom must be used for hand hygiene (no other option available), avoid contamination of hands with potentially contaminated surfaces and objects.

Bar soaps are not acceptable in healthcare facilities or community settings for use by HCWs, and are only for individual P/R/C use. In healthcare facilities bar soap should be supplied in small, single P/R/C use pieces, and must be stored in a soap rack to allow drainage and drying. Discard bar soap on discharge.

Plain soaps act on hands by emulsifying dirt and organic substances (e.g., blood, mucous), which are then rinsed away with running water. Antimicrobial agents in plain soaps are only present as a preservative.^{13.6}

- Antimicrobial soaps have residual antimicrobial activity and are not affected by the presence of organic material.

Disadvantages of antimicrobial soap include:

- Antimicrobial soaps are harsher on hands than plain soaps and frequent use may result in skin breakdown; and
- Frequent use of antimicrobial soap may lead to antibiotic resistance in microorganisms.

Hand hygiene with correctly applied ABHR kills microorganisms in seconds; Hand hygiene with soap and water done correctly, physically removes microorganisms

2.3. Care Environments

The care environment is the space around the P/R/C that may be touched by either that P/R/C or staff.

Two different environments:

1. Healthcare External Environment/Zone:

This is the environment beyond the immediate area surrounding the P/R/C.

- In a single bed room this is outside the room
- In a multi-bed room this is everything outside the bed area of the P/R/C
- In home care this is equipment and transport or storage containers temporarily brought in to the home.^{13.2} (examples include: supply/visit carry bag, equipment, pens etc. – if items have to be brought into P/R/C environment clean/disinfect immediately before and/or after use)
- This includes the people within it; staff, visitors, volunteers and other P/R/C are part of the healthcare (external) environment. In the home this would include other household members ^{13.2}

NOTE: For staff this means their uniform/pockets, glasses, hair, etc. are part of the healthcare (external) environment.

2. Patient/Resident/Client (P/R/C) Environment/Zone:

The term 'patient/resident/client (P/R/C) zone' refers to the space that contains the P/R/C, as well as the immediate surroundings and inanimate surfaces in contact with that P/R/C (e.g., bed rails, chair, bedside tables, work surfaces, bed linens, infusion tubing, and other medical equipment). It also contains surfaces frequently touched by staff within the vicinity of the P/R/C (e.g., monitors, buttons and knobs, and other frequently touched - "high touch" surfaces within the P/R/C zone).

- In a single room this is everything in the room of the P/R/C

- In a multi-bed room this is the area inside the privacy curtain/divider space of the P/R/C
- In an Emergency department cubicle, it is the stretcher of the patient and the equipment in close proximity used in the care of the patient
- In a nursery/neonatal and intermediate care setting, the patient environment includes the inside of the bassinette or isolette, the equipment outside the bassinette or isolette used for that infant (e.g., ventilator, monitor), as well as an area around the infant (i.e., within approximately 1 metre/ 3 feet)
- In an ambulatory care/clinic setting this is the P/R/C themselves, their belongings and any equipment/furniture being used during care/treatment
- In home care this is the entire residence of the P/R/C^{13.2}

2.4. Indications and Moments for Hand Hygiene During Healthcare Activities

When shall hand hygiene be performed? A hand hygiene indication identifies why hand hygiene is necessary at a given moment. There may be several indications to perform hand hygiene in a single care sequence or activity. Hand hygiene shall be performed before and after any direct contact with a P/R/C or their equipment, between procedures on the same P/R/C, and before contact with another P/R/C. While all indications for hand hygiene are important, there are some essential moments in healthcare settings where the risk of transmission is greatest and hand hygiene must be performed.

Essential HH indications can be simplified into 4 moments for training

3. THE 4 MOMENTS OF HYGIENE

3.1. MOMENT 1: BEFORE INITIAL PATIENT/RESIDENT/CLIENT (P/R/C) CONTACT OR P/R/C ENVIRONMENT CONTACT

WHEN? *Clean your hands when entering a P/R/C environment*

Examples include but are not limited to:

- **Before entering** the P/R/C room/bed-space/home, treatment/exam room
- **Before touching** P/R/C (e.g., shaking their hand, helping them move around)
- **Before touching any object** or furniture in the P/R/C's environment (e.g., stretchers, wheelchairs, infusion rate adjustment, silencing a pump).

WHY? *To protect the P/R/C and their environment from harmful microorganisms (germs) carried on your hands*

3.2. MOMENT 2: BEFORE ASEPTIC/CLEAN PROCEDURES

WHEN? *Clean your hands immediately before any aseptic procedure*

Examples include but are not limited to:

- Performing invasive procedures
- Handling dressings or touching open wounds
- Preparing and administering medications
- Preparing, handling, serving or eating food
- Feeding a P/R/C
- Before and after shifts and breaks

WHY? *To protect the patient/resident/client from harmful microorganisms, including a P/R/C's own microorganisms, entering his or her body*

Consideration should be given to communal staff rooms and foods. ABHR should be available and items should be individually packaged or distributed in a controlled manner (i.e., supervised, using tongs, appropriate hand hygiene, etc.)

3.3. MOMENT 3: AFTER BODY FLUID EXPOSURE RISK

WHEN? *Clean your hands immediately after an exposure risk to blood and body fluids, non-intact skin, and/or mucous membranes (and after glove removal)*

Examples include but are not limited to:

- Contact with blood and body fluids
- Contact with items known or considered to be contaminated
- Procedures on the same P/R/C where soiling of hands is likely, to avoid cross-contamination of body sites
- Oral care, wound care, patient/resident/client toileting
- Removal of gloves
- Personal use of toilet or wiping nose/face
- Feeding a P/R/C
- Before and after shifts and breaks.

WHY? *To protect yourself and the healthcare (external) environment from harmful patient/resident/client microorganisms*

3.4. MOMENT 4: AFTER PATIENT/RESIDENT/CLIENT CONTACT OR PATIENT/RESIDENT/CLIENT ENVIRONMENT CONTACT

When? *Clean your hands when leaving the P/R/C, and/or P/R/C environment*

Examples include but are not limited to:

- **After touching P/R/C** to assist with any tasks (e.g., helping a patient/resident/client mobilize; giving a massage; taking a pulse, blood pressure, chest auscultation, abdominal palpation)
- After touching any object or furniture in the P/R/C's environment (e.g., changing bed linen, infusion rate adjustment, alarm monitoring, clearing the bedside or overbed table)
- When leaving a patient/resident/client's home.

Why? *To protect yourself and the healthcare (external) environment from harmful patient/resident/client microorganisms*

Hand hygiene with point of care alcohol based hand rub (ABHR) is the standard of care expected of all staff, in all healthcare settings. Busy staff need access to hand hygiene products anywhere care is provided to a P/R/C or contact with their environment is taking place (from the ICU to the community outreach clinic). Making ABHR available at the point of care (e.g., within arm's reach) is an important system support to improve hand hygiene. This enables staff to quickly and easily follow the [4 Moments for Hand Hygiene](#).

Point of care ABHR can be achieved with a variety of methods (e.g., ABHR attached to the bed, wall, containers carried by the healthcare worker [HCW]). Sites/programs shall complete a risk assessment to determine the most appropriate placement of ABHR in every facility and provide ABHR at point-of-care so it is easily accessible for appropriate use. To avoid confusion, ABHR dispensers should not be placed near hand washing sinks.

Two moments for hand hygiene may sometimes fall together. Typically, this occurs when going from one P/R/C to another without touching any surfaces when moving from one P/C/R zone to another. Naturally, a single hand hygiene action will cover the two moments for hand hygiene.

For example: Performing HH after touching a P/R/C ([Moment 4](#)) would also cover doing HH before touching another P/R/C ([Moment 1](#)). Two Moments are covered by performing HH once.

3.5. HAND HYGIENE TECHNIQUES

- Remove hand and arm jewelry as these items are hard to clean and prevent the removal of microorganisms from surfaces of the hands and wrists they cover [13.3](#)
- If a watch is worn, it must be worn above the wrist and fit snugly
- Avoid long sleeves. Clothing or other items that impede frequent and effective hand hygiene should be removed.

3.5.1. Using an Alcohol-Based Hand Rub (ABHR)

- Ensure hands are visibly clean (if soiled, follow hand washing steps)
- Apply one to two full pumps of product (about 1.1 - 2.0 ml) onto one palm; the volume should be enough so that 15 seconds of rubbing is required for drying ([13.12](#))
- Rub product over all surfaces of hands, concentrating on finger tips, between fingers, back of hands, wrists and base of thumbs; these are the most commonly missed areas; and continue rubbing hands until product is dry; this will take a minimum of 15 seconds if sufficient product is used. Hands must be fully dry before touching the P/R/C or the care environment/equipment for the ABHR to be effective. This also eliminates the extremely rare risk of flammability in the presence of an oxygen-enriched environment. DO NOT WIPE OFF.
- There is insufficient evidence for the efficacy of non-alcoholic, waterless antiseptic agents in the health care environment. Therefore, they shall not be used in health care settings.

3.5.2. Using Soap and Water

- Wet hands with warm (not hot or cold) water; hot or cold water does not significantly impact microbe removal but it is hard on the hands, and will lead to dryness
- Apply liquid/gel or foam soap
- Vigorously lather all surfaces of hands for a minimum of 15 seconds to create a good lather; removal of transient or acquired microorganisms
- Microorganisms require a minimum of 15 seconds of mechanical action; Pay particular attention to finger tips, between fingers, backs of hands, wrists and base of the thumbs; these are the most commonly missed areas
- Using a rubbing motion, thoroughly rinse soap from hands with running water; residual soap can lead to dryness and cracking of skin
- Dry hands thoroughly by blotting hands gently with a paper towel; rubbing vigorously with paper towels can damage the skin

- Turn off taps with paper towel to avoid recontamination of the hands. If hand air dryers are used in non-clinical areas, hands-free taps are required
- DO NOT use ABHR immediately after washing hands, as skin irritation will be increased
- HH should not be performed at a P/R/C's sink as this may re-contaminate hands. HH should be done as soon as a dedicated HH sink is available
- If the P/R/C bathroom must be used for hand hygiene (no other option available), avoid contamination of hands with potentially contaminated surfaces and objects.

IF performing hand hygiene with soap and water:

- The sink and the area surrounding the sink must be visibly clean and running water must be available
- The sink should be at point of care
- Use only liquid/gel or foam soap to wash and paper towels to dry to your hands. The sink should be at point of care
- If any of these things are not available ABHR must be used.

3.5.3. When Hands are Visibly Soiled and Liquid/Gel Soap and Running Water is Not Available

- Use a moist pre-packaged wipe to remove the soiling and then follow with ABHR to perform hand hygiene
- Wash hands once a suitable sink and hand hygiene supplies are available.

3.6. Factors that Reduce Effectiveness of Hand Hygiene

3.6.1. Condition of the Hands

The condition of the hands can influence the effectiveness of hand hygiene. Intact skin is the body's first line of defense against microorganisms; therefore, hand care is an essential part of the hand hygiene program. See section [3.6.9 – Lotions and Creams](#) for more information. The presence of dermatitis, cracks, cuts or abrasions can trap microorganisms and compromise hand hygiene. Dermatitis also increases shedding of skin squamous (cells) and, therefore, shedding of microorganisms. If there are any concerns regarding skin integrity, consult Occupation and Environmental Safety and Health (OESH).

3.6.2.Nails

Long nails are difficult to clean, can pierce gloves and harbour more microorganisms than short nails. Keep natural nails clean and short. The nail should not show past the end of the finger. Clean, short fingernails (no more than 0.64 cm or ¼ inch) are required by direct care staff that comes into contact with:

- | | |
|---|--|
| <input checked="" type="checkbox"/> Food | <input checked="" type="checkbox"/> Sterile linens/supplies |
| <input checked="" type="checkbox"/> Equipment used for care | <input checked="" type="checkbox"/> Patients/Residents/Clients |
| <input checked="" type="checkbox"/> Blood or body fluids | <input checked="" type="checkbox"/> The care environment |

3.6.3.Nail Polish

- Nail polish, if worn, must be fresh and in good condition (i.e., not chipped)[13.4](#)
- Nail polish cannot be worn for more than 4 days and must be removed when it becomes chipped
- Nail polish that is chipped or worn longer than four days can harbor microorganisms that are not removed by hand washing, even with surgical hand scrubs[13.5](#)
- Freshly applied nail polish does not result in increased numbers of bacteria around the nails
- Gel polish has been shown to damage nails, resulting in nail weakness, brittleness and thinning, putting nails at increased risk for breaking. Nail art (adding decorative paint effects to nails) has been shown to be associated with outbreaks of infection.[13.4](#)

3.6.4.Artificial Nails or Nail Enhancements

Artificial nails and nail enhancements (gel nails, wraps or extenders - adhesive decorative plastic or vinyl attached to nails) are not to be worn by direct care staff (those who come into contact with food, equipment used for care, blood or body fluids, sterile linens/supplies, P/R/Cs and the care environment. Refer to [WRHA Policy 20.10.020, Dress Code](#), and [WRHA Policy 90.00.060 Routine Practices for Reducing the Risk of Infection Transmission](#).

Artificial nails harbor more microorganisms and are more difficult to clean than natural nails. Artificial nails and nail enhancements have been implicated in the spread of microorganisms and in outbreaks, particularly in neonatal nurseries and other critical care areas. Surgical site infections and hemodialysis-related bacteremias have also been linked to artificial nails. Artificial nails and nail enhancements are also associated with poor hand hygiene practices and result in more tears to gloves.[13.4](#)

3.6.5. Jewelry

Hand and arm jewelry hinder hand hygiene. Rings increase the number of microorganisms present on hands and increase the risk of tears in gloves. Direct care staff are encouraged to remove hand and arm jewelry prior to work. These items are hard to clean and prevent the removal of microorganisms from surfaces of the hands and wrists that they cover.[13.6](#) A simple and practical solution allowing effective hand hygiene is for HCWs to wear their rings around their neck on a chain as a pendant.[13.6](#)

If watches and other wrist jewelry are present, remove or push up above the wrist before performing hand hygiene. They should not interfere with or become wet during hand hygiene. In areas where earrings must be removed or covered with PPE, facial jewelry shall be treated the same way as pierced earrings, i.e., staff must remove or confine all facial jewelry when in areas where pierced earrings must be removed or covered with PPE.

3.6.6. Upper Extremity Support Devices

Direct care staff (those that come into contact with what is listed in [3.6.2 Nails](#)) who wear an upper extremity support device (UESD) must be able to clean the device and perform hand hygiene as per the direction in the [Upper Extremity Supportive Devices Infection Prevention and Control Communication Form](#). See [OESH Upper Extremity Support Devices Operational Directives](#) if more information is needed.

3.6.7. Other Obstacles to Effective Hand Hygiene

- Long sleeves should not interfere with, or become wet, when performing hand hygiene
- Ensure long hair is tied up and off the collar to avoid inadvertently touching hair following a hand hygiene moment.

Missed opportunities (seen in hand hygiene audits) are also observed when staff touch their own clothing, personal items, face or equipment (e.g., stethoscope, coffee cup, chart, etc.)

3.6.8. Hand Drying (paper towel, air dryers)

Effective hand drying is important for maintaining hand health. Considerations include:

- Disposable paper hand towels provide the lowest risk of cross-contamination and should be used for drying hands in clinical practice areas (e.g.: P/R/C rooms, clinic rooms etc.)
- If cloth drying towels are used, a new towel must be used for each hand hygiene episode
- Towel dispensers must be mounted so access to them is unobstructed and splashing or dripping onto adjacent wall and floor surfaces is minimized
- Towel dispenser design should be designed so only the towel is touched during its removal for use
 - Towels hanging from the dispenser should not hang directly into a garbage can
- Hot-air dryers, including jet air dryers, must not be used in clinical areas as warm air currents dry hands slowly and can be used by only one person at a time. This results in lines and the temptation to dry hands on clothing. Germs are drawn into the hot-air dryers and redeposited onto freshly washed hands. These germs are also recirculated into the air.

3.6.9. Lotions and Creams

- To be effective, skin care products should be used regularly. Health care facilities/programs should develop a proactive program to keep hands healthy so hand hygiene can be optimal
- HCWs must use facility approved lotions compatible with products and gloves in use
- Position skin care products as close as possible to areas where hand hygiene is performed
- Use dispensers of sufficient quality that they will not clog or leak
- Hand lotion bottles shall not be reused
- Barrier Creams: unlike hand lotions, which penetrate the skin via pores, barrier creams are adsorbed to the skin and are designed to form a protective layer that is not removed by standard hand washing. Barrier creams may actually be harmful as they trap agents beneath them, ultimately increasing risk for either irritant or allergic contact dermatitis
- Inappropriate barrier cream application on HCW hands may exacerbate irritation rather than provide benefit.

3.6.10. Dispensers

- Products must be dispensed in a disposable pump/squirt container that is not topped-up, to prevent contamination
- **DO NOT** add soap or hand rub to a partially empty dispenser
- If reusable dispensers/containers are utilized the container as well as the pump system must be emptied, washed and air-dried completely prior to refilling
- Locked, tamper-proof containers should be used to secure the product in place
- National Fire Code, and local fire regulations, for ABHR placement and storage shall be adhered to. Consideration of alternate ABHR options may be required to support necessary point-of-care use, as well as adhere to fire regulations
- An environmental risk assessment should be performed to determine the most appropriate placement of ABHR dispensers. [13.1](#)

Careful selection of products used for hand hygiene practice (e.g., ABHR, soaps, lotions, paper towels) has a significant impact on hand hygiene compliance

4. SOURCE CONTROL

These measures are used to minimize the spread of microorganisms (germs) from an infectious source. Symptomatic persons require direction at the point of initial encounter and in strategic places in any healthcare setting to minimize potential infectious spread (e.g., triage, reception and waiting areas, elevators, cafeterias).

Source control measures may include but are not limited to:

- Signage at healthcare setting entrances for early identification of symptoms
- Hand hygiene
- Separate entrances/waiting areas for persons with a potential infection
- Spatial separation
- Physical barriers for acute assessment
- Early identification, diagnosis and treatment of infection
- Respiratory etiquette/hygiene
- Placement of the P/R/C requiring Additional Precautions (e.g.: single rooms/airborne infection isolation rooms [AIIRs]).

4.1. Respiratory Etiquette/Respiratory Hygiene

Respiratory hygiene refers to a combination of measures designed to decrease the spread of respiratory microorganisms. These 'source control' measures are targeted to all persons with symptoms of respiratory infection throughout every encounter in the healthcare setting.

Respiratory hygiene involves educating and encouraging everyone (P/R/Cs, HCWs designated caregivers and visitors) who have the physical and cognitive abilities to do so, to practice respiratory hygiene. Specific measures may include instructional signs, education programs and provision of materials for respiratory hygiene (e.g., tissues, plastic lined waste receptacles, alcohol-based hand rub [ABHR], medical masks).

Respiratory Hygiene includes:

- Covering your mouth and nose against your sleeve/shoulder during coughing or sneezing
- Using tissues to contain respiratory secretions by covering your mouth and nose during coughing or sneezing, with prompt disposal of the tissue into a hands-free garbage
- Wearing a medical mask when coughing or sneezing
- Turning your head away from others when coughing or sneezing
- Maintaining a spatial separation of two metres/six feet between persons that are symptomatic with an acute respiratory infection and those who do not have symptoms of a respiratory infection. If this cannot be achieved, the person with respiratory symptoms must be at least one metre/three feet apart and the symptomatic person must wear a medical mask. One metre/three feet may be sufficient for young children and others whose cough is not forceful enough to propel the droplets as far as two metres/six feet.

**Family/visitors with signs/symptoms of respiratory illness SHOULD NOT visit.
Staff with sign/symptoms of an illness should stay home**

4.2. Triage

4.2.1. Emergency Rooms and Acute Assessment Settings

- Post signs to direct persons with symptoms of acute infection (e.g., cough, fever, vomiting, diarrhea, coryza (nasal congestion), rash, and conjunctivitis) to specific waiting areas
- Ensure a physical barrier (e.g., plastic partition at triage desk, wall, portable wipeable divider) is located between infectious sources (e.g., those with symptoms of a respiratory infection) and others

- Place P/R/Cs who are likely to contaminate the environment directly into a single examination room whenever possible. For example, P/R/Cs with:
 - Gastrointestinal (acute diarrhea/vomiting) illness
 - Respiratory infections. These persons should be placed either directly into an examination room or an airborne infection isolation room, as indicated by the respiratory infection suspected. **Place a medical mask on these persons until isolated or spatial separation is achieved**
 - Excessive bleeding or body fluid drainage into a single examination room whenever possible.

4.2.2. Ambulatory Care/Clinic Settings

- If possible, identify persons with symptoms of an acute infection when scheduling appointments for routine clinic visits and request, if possible, then defer routine clinic visits until symptoms of the acute infection have subsided
- Inform those who cannot defer their routine clinic visit (i.e., those that require assessment of symptoms/condition) to follow hand hygiene and/or respiratory hygiene recommendations appropriate for their symptoms. Direct these persons into an examination room as soon as they arrive and/or schedule their appointment for a time when other persons seeking care are not present
- Post signs at clinic entrances reminding symptomatic persons to perform [hand hygiene](#) and/or [respiratory hygiene](#) if they have symptoms.

4.3. Early Diagnosis and Treatment

Ensure symptomatic P/R/Cs receiving care are assessed in a timely manner and that potential communicable infection(s) are considered (e.g., tuberculosis, norovirus, respiratory syncytial virus [RSV], pertussis).

4.4. Spatial Separation

Appropriate spatial separation and spacing requirements are necessary to decrease exposure to microorganisms (germs) for everyone in clinical and waiting areas. There should be a two metres/six feet spatial distance between a coughing/sneezing infected source (e.g., symptomatic P/R/Cs with acute respiratory infection with a cough, fever or shortness of breath) and an unprotected susceptible host (e.g., P/R/C, HCWs, visitors, contractors). This is recommended to prevent the transmission of droplet borne infectious particles. In inpatient/resident facilities, a single room with in-room designated toilet and sink is preferable, as it may be difficult to maintain the recommended spatial separation of two metres/six feet between P/R/Cs.

If two metres/six feet cannot be achieved, those receiving care must be at least one metre/three feet apart and the symptomatic P/R/C must wear a medical mask. Always

ensure the medical mask covers the mouth and nose. One metre/three feet may be sufficient for young children and others whose cough is not forceful enough to propel the droplets as far as two metres/six feet.

4.5. Aerosol-Generating Medical Procedures (AGMPs)

Aerosol-generating medical procedures can generate aerosols as a result of artificial manipulation of a person's airway. Several types of AGMPs have been associated with an increased risk of tuberculosis (TB), Severe Acute Respiratory Syndrome (SARS), and Middle Eastern Respiratory Syndrome (MERS CoV) transmission. While there is some evidence for the spread of infections via droplets and aerosols by these procedures, further research is needed to quantify the risk. Infection transmission may increase during AGMPs because of the potential to generate a high volume of respiratory aerosols that may be propelled over a longer distance than with natural dispersion. These procedures include:

- ☑ Endotracheal intubation and extubation, manual bag mask ventilation, insertion of laryngeal mask airway (LMA)^{13.7}
- ☑ Bronchoscopy and bronchoalveolar lavage
- ☑ Tracheostomy procedure (open or percutaneous) Laryngoscopy (with instrumentation below the vocal cords)
- ☑ Non-invasive positive pressure ventilation (BiPAP and CPAP) High flow nasal cannula oxygenation (e.g. Optiflow) - should only be used in patients with COVID-19 following consultation with an Attending Intensivist
- ☑ Open deep suctioning via endotracheal tube/tracheostomy
- ☑ Cardiopulmonary resuscitation (with manipulation of the airway)
- ☑ Sputum induction using hypertonic saline
- ☑ Some dental procedures (e.g., high speed drilling, ultrasonic scalers etc.)
- ☑ Autopsy of lung tissue
- ☑ Administration of nebulizing medications, does not include administration of a metered dose inhaler (MDI).

The following are NOT considered AGMPs:

- Oxygen delivered via nasal prongs and/or non-rebreathe masks are not considered AGMP, regardless of flow rate.
- Chest compressions are not considered an AGMP
- Collection of nasopharyngeal swabs and/or nasopharyngeal aspirates are not considered AGMPs, there is no published literature documenting transmission of respiratory infections, including TB, SARS, influenza, and COVID-19 by collection of these specimens.

**** Specific COVID AGMP** guidance can be found at: [aerosol-generating-medical-procedures-AGMPs.pdf \(sharedhealthmb.ca\)](https://sharedhealthmb.ca/aerosol-generating-medical-procedures-AGMPs.pdf) **

Routine Practices are sufficient for AGMPs performed on P/R/C with no signs or symptoms of suspected or confirmed TB, SARS, COVID or respiratory infections due to an emerging respiratory pathogen^{13.10}

Prior to performing any AGMP, P/R/C should be carefully assessed and [Strategies to Reduce Risk from AGMPs](#) should be followed when the P/R/C is suspected or confirmed as having any of the following:

- Airborne Pathogens (i.e.: tuberculosis)
- SARS
- MERS CoV
- SARI
- COVID
- Viral Hemorrhagic Fever (VHF)
- Or other emerging respiratory infections

STRATEGIES TO REDUCE RISK OF AGMPs

1. Carefully analyze risks and benefits to AGMPs; avoid performing unnecessary AGMPs.
2. Consider alternative to AGMPs.
3. Anticipate and plan for AGMPs, including using appropriate engineering controls (airborne infection isolation rooms or private rooms, evaluating air exchange rates, personal protective equipment, etc.).
4. Depending on the procedure, sedation may be appropriate for the P/R/C requiring the AGMP, to minimize excessive and/or prolonged and/or forceful coughing etc.
5. Paralytics to minimize the risk of aerosolization (for intubation or if the patient's breathing is already supported by mechanical ventilation) can be used when appropriate.
6. Use closed endotracheal suction systems whenever possible.
7. Use the minimum required number of staff in the room when performing an AGMP.
8. Ensure appropriate PPE is worn by all staff present in the room during the procedure. PPE guidance can be found in the Provincial Guidance for Aerosol Generating Medical Procedures (AGMPs) for COVID.
9. Choose an appropriate space for an AGMP. The appropriate space for an AGMP will vary depending on the patient and the circumstances in which the AGMP is taking place.
10. Once an AGMP is complete make sure the door to the room remains closed and staff continue to wear N95s until appropriate air exchanges (at 99% minimally, and ideally at 99.9%) have occurred according to the [Air Exchanges Table](#).

NOTE: *When responding to a code (e.g., cardiac arrest) for a P/R/C requiring airborne isolation, when neither an AIIR or single room with door closed are available for an AGMP, draw the privacy curtains and remove any shared equipment, supplies or linens from the immediate vicinity prior to performing an AGMP. Ensure all staff are wearing appropriate personal protective equipment and remove everyone else in the room where possible.*

5. ACCOMMODATION AND PLACEMENT

Accommodation of patients/residents/clients (P/R/C) in single rooms improves infection prevention and control. Single rooms with a private toilet, designated hand washing sink for P/R/Cs, and designated staff hand washing sink may reduce opportunities for cross transmission particularly when the P/R/C has poor hygiene, contaminates the environment or cannot follow IP&C measures because of age or decreased cognitive abilities.

5.1. Options for P/R/C Placement and Room Sharing

If the availability of single rooms is limited, use the [Point of Care Risk Assessment](#).

Consider if the P/C/R:

- Has presence or absence of known or suspected infection and its route(s) of transmission (i.e., need for [Additional Precautions](#)).
 - [Contact Precautions](#) (single room is preferred)
 - [Droplet Precautions](#) (single room is preferred)
 - [Airborne Precautions](#) (airborne infection isolation room [AIIR] required).
- Visibly soils the environment or they cannot maintain appropriate toileting and respiratory hygiene
- Has uncontained secretions or excretions
- Has wound drainage that cannot be contained by a dressing
- Has fecal incontinence where stools cannot be contained in incontinent products or infant diapers.

5.2. Risk Factors for Transmission from the Infected P/R/C?

- Are roommates susceptible to adverse outcome from a healthcare associated infection (HAI)?
- Are there options for room sharing (e.g., cohorting P/R/Cs infected with the same organism)?

IF SO...

- Can the P/R/C's roommate(s) and visitors follow infection prevention and control measures?
- Give priority for placement in single rooms to those who pose an increased risk for transmission of a microorganism to others
- If AIIRs are in limited supply/high demand, refer to Priority of AIIRs.

5.3. Priority for Single Rooms Goes to Those:

- Needing Additional Precautions
- Identified as high risk for transmission of microorganisms (e.g., stool incontinence, uncontained secretions)
- Identified as being at higher risk of acquisition and adverse outcomes resulting from transmission of microorganisms (e.g., immunosuppression, open wounds, indwelling catheters, and anticipated long length of stay)
- Requiring dependence on staff for activities of daily living.

5.4. Factors to be considered with shared rooms include:

- The selection of appropriate roommates
- Avoid placing P/R/C at high risk of complications, if they should become infected, in rooms with P/R/C with transmissible infections, diarrhea or open wounds
- Clearly define the boundary of the potentially contaminated P/R/C area within the shared room (e.g., draw privacy curtain/place portable divider around P/R/C)
- Prevent transmission risks through avoiding the sharing of sinks and toilets
- Assessing activities of the roommates and their visitors (e.g.: is the P/R/C a wanderer, will visitors follow the correct precautions if interacting with the person on contact precautions, etc.).

5.5. Cohorting

Assignment of P/R/Cs known to be infected with the same microorganisms/strain to the same room (cohorting) or separate units or areas has been successful in controlling transmission of some microorganisms. Contact IP&C to determine appropriate cohorting.

5.6. The Use of Airborne Isolation Rooms (AIIRs)

AIIRs are designed with negative pressure ventilation (i.e., with air flow from the outside corridor into a room through the doorway and exiting directly to the outside of the building or filtered before recirculation). They are used for accommodation of P/R/C suspected or confirmed to have an infection spread by the airborne transmission route.

An AIIR is also required when performing AGMPs on those with SARS, MERS CoV, COVID, viral hemorrhagic fever and other emerging pathogens for which transmission characteristics are not yet known.

In settings where AIIRs are limited, the following process should be used to assess the accommodation and/or continued accommodation along with clinical judgement and risk/benefit analysis. This will be used to determine the risk of infectivity and risk of transmission and/or disease and exposure to others. This risk assessment should be done

in collaboration with IP&C/designate, Public Health/delegate and other key staff involved with the care.

Factors to be included in the risk assessment for an AIIR (done with the Infection Control Professional/designate), but not limited to, are:

- Degree of transmissibility of the infectious disease
- Presence of communicable symptoms (e.g., coughing)
- Potential and level of the P/R/Cs infectivity
- Stage of recovery of the P/R/C
- Immune status of others.

In situations when AIIRs are not available, conduct a risk assessment looking at the factors identified above. The P/R/C can be temporarily housed in a single room with the door closed, away from high risk persons. P/R/C requiring an AIIR should be transferred as soon as medically feasible to a facility/unit with AIIRs. If AIIRs in other facilities are not available, a decision should be made following the risk assessment above to determine if it will be safe to accommodate and/or treat the P/R/C in the facility and whether or not that P/R/C should continue to be masked while in the room.

Home Settings

Individuals who have not been exposed or are not immune should be advised to avoid sharing airspace with the client requiring precautions. Natural ventilation (e.g., open windows) will help disperse the microorganisms from the room. Advise the client to exclude themselves from group programs, routine services that are not medically necessary (e.g.: interactions with volunteers) when experiencing acute symptoms of an infection.

6. FLOW

Flow refers the transfer and transport of the patient/resident/client (P/R/C) ***within and outside*** of the facility. There is a potential for exposure to, and spread of microorganisms (germs), as a result of the activity or transport of the P/R/C due to unintended contact with others, items used for care, and environmental surfaces.

Patients/Residents/Clients should not be transported between units, departments or facilities unless medically necessary.

Frequent transfers should be avoided as this increases the number of interactions with staff and others, and provides opportunities for transmission to occur. Staff, including bed/accommodation coordinators, are responsible for selecting the most appropriate accommodation based on the [PCRA](#) and for prioritizing use of single rooms and AIIRs if they

are limited. Using the [PCRA](#) can minimize unnecessary transfers. When in doubt regarding transfers and accommodation, consult IP&C/designate.

6.1. Flow and Additional Precautions

- Advance communication between the transporting area and the receiving area is important to ensure precautions are used correctly and to decrease unnecessary waiting time in public areas
- Use source control measures (e.g., request that the P/R/C being transported/ transferred perform [hand hygiene](#) before leaving their room, cover skin lesions, wear clean clothes, wear a mask, etc.).

6.2. Ambulatory Care/Clinic Setting

When Additional Precautions are necessary, those scheduled for an appointment should defer when possible (e.g., routine foot care) or enter through a separate entrance. Upon arrival, P/R/Cs requiring Additional Precautions should be asked to perform [hand hygiene](#), apply [PPE](#) if appropriate (e.g., medical mask), and be placed in an examination room. The door of the exam room should be closed if an airborne spread microorganism is suspected (e.g., measles, tuberculosis).

7. ASEPTIC TECHNIQUE

Aseptic technique, sometimes referred to as sterile technique, means using practices and procedures to prevent contamination from microorganisms (germs) on the patient/resident/client (P/R/C) skin or another person's flora to a sterile body site. These practices are required when performing procedures that expose the person's normally sterile body sites (e.g., intravascular system, spinal canal, subdural space, urinary tract). Practices such as creating a sterile field or preparing skin with an antiseptic significantly reduces the risk of introducing microorganisms that can lead to infections.

Components of aseptic technique prior to a procedure may involve the following:

- Preparing the P/R/C's skin with an antiseptic
- [Hand hygiene](#), preferably with ABHR, or if not accessible, an antimicrobial soap if it is an invasive procedure (e.g. placing central intravascular catheters or catheters for injecting into the spinal canal or subdural spaces)^{13.1}
- Sterile gloves
- Gowns
- Medical masks (procedure or surgical masks), where required, to prevent microorganisms carried in the HCW's nose and mouth from contaminating the sterile field
- Sterile drapes, used to prevent transferring microorganisms from the environment to the P/R/C while the procedure is being performed
- Maintaining a sterile field.

Infections may result from failure to use proper skin antisepsis prior to injection of medications, vaccines or venipuncture.

Chlorhexidine in alcohol inactivates microorganisms on the skin more effectively than most other antiseptics and is the preferred antiseptic for skin preparation prior to insertion of central venous catheters and pulmonary artery catheters.

7.1. Recommendations for Injection Safety Include

- Perform [hand hygiene](#) prior to accessing supplies, handling vials and IV solutions, and preparing or administering medications [13.4](#)
- Use aseptic technique in all aspects of parenteral medication administration, medication vial use, injections and glucose monitoring procedures. Limit access to select trained persons, if possible
- **NEVER** administer medications from the same syringe to more than one P/R/C, even if the needle is changed
- Consider a syringe or needle contaminated after it has been used to enter or connect to a P/R/C's intravenous infusion bag or administration set
- **DO NOT** enter a vial, bag or bottle with a syringe or needle that has been previously used
- **NEVER** store needles and syringes unwrapped as sterility cannot be assured [13.3](#)
- Assign medications packaged as multi-use vials to a single P/R/C whenever possible
- **DO NOT** use bags or bottles of intravenous solution as a common source of supply for more than one patient/resident/client
- Provide a puncture resistant sharps container that is available at point of use [13.4](#)
- Store and prepare medications and supplies in a clean area on a clean surface [13.4](#)
 - Label sterile solutions containers with the date opened and discarded every 24 hours and/or according to manufacturer's instructions [13.4](#)
 - Discard outdated medications. There should be a process in place to check expiry dates before use. [13.4](#)

7.2. Aseptic Technique for Invasive Procedures and Handling Injectable Products

- Perform [hand hygiene](#), with ABHR prior to opening supplies:
 - When ABHR is not accessible, perform hand hygiene with antimicrobial soap and water.
- Open tray and supplies only when ready to use to ensure a sterile field
- Perform [hand hygiene](#) prior to applying personal protective equipment, as indicated by the specific procedure

- Prepare the skin of the P/R/C with an appropriate antiseptic before performing an invasive procedure
- Use the appropriate size drape when a drape is required, to maintain a sterile field.
- **DO NOT** administer medications or solutions from single dose vials, ampules or syringes to multiple P/R/C s or combine leftover contents for later use
- Use a sterile, single use disposable needle and syringe for each medication/fluid withdrawal from vials or ampules
- Clean and disinfect the stoppers or injection ports of medication vials, infusion bags, etc., with alcohol before entering the port, vial or bag
- **Use single dose medication vials, prefilled syringes, and ampules in clinical settings.** If the product is only available as multi-dose vials, [see multi-dose vials](#) below.

7.3. Single Dose Vials

Single Dose vials, intended for single patient/resident/client use, typically lack preservatives. The use of these vials for multiple P/R/Cs carries a substantial risk for bacterial contamination.

- Use single dose medication vials, prefilled syringes, and ampules in clinical settings. If the product is only available as multi-dose vials, [see multi-dose vials](#) below
- **NEVER** use medications packaged as single use vials for more than one P/R/C
- **ALWAYS** use a sterile syringe and needle/ cannula when entering a vial.
- **NEVER** enter a vial with a syringe or needle/cannula that has been used on a P/R/C.

7.4. Multi-Dose Vials

Transmission of hepatitis B and hepatitis C has followed the reuse of needles and/or syringes when withdrawing from multi-use vials.

- Restrict the multi-dose vial to single P/R/C use whenever possible
- A multi-dose vials should only be used when a product is only available for purchase in multi-dose vials
- Prepare syringes from multi-dose vials from a centralized medication preparation area (i.e., do not take multi-dose vials to the P/R/C environment)
- Store the multi-dose vial in a restricted access location (e.g., in a secure location away from P/R/C bedside and where access is restricted, such as a medication room or locked cart)
- Cleanse the access diaphragm of vials using friction and 70% alcohol. Allow to dry before inserting a needle into the vial^{13.3}

- Use a sterile, single use needle and syringe each time the multi-dose vial is entered
 - **DO NOT** re-enter the multi-dose vial with a previously used needle or syringe
- Label the multi-dose vial with date of first opening. See the product manufacturer's instructions for use for recommended durations of use after entry of a multi-dose vial [13.3](#)
- Discard opened multi-dose medication vials according to the manufacturer's instructions or 28 days after opening, whichever is shorter [13.3](#)
- Inspect the multi-dose vial for clouding or particulate contamination prior to each use and discard multi-dose vial if clouding or particulate contamination present
- Discard the multi-dose vial if sterility or product integrity is compromised
- **NEVER** leave a needle in a multi-dose vial. [13.3](#)

7.5. Single P/R/C Multi-Use Devices

Assign single P/R/C multi-use devices (e.g., glucose sampling devices, finger stick capillary blood sampling devices) to only one P/R/C. If it is not feasible to assign glucose meters to one P/R/C, clean and disinfect before use on others.

7.6. Injecting Material and Placing a Catheter into the Spinal Canal or Subdural Space

Use [aseptic technique](#) including creating a sterile field, aseptic skin preparation and use of a medical mask and sterile gloves (e.g., during lumbar puncture, myelogram, and spinal or epidural anesthesia).

7.7. Insertion of Central Venous Catheters

- Use maximal aseptic barriers as outlined in [7.2 Aseptic Technique for Invasive Procedures and Handling Injectable Products](#) (above), in addition to a cap, medical mask, long sleeved sterile surgical gown, sterile gloves, and a large full body sterile drape
- Prepare the skin with chlorhexidine in alcohol or an equal alternative for inserting any central venous catheter or pulmonary catheter.

7.8. Insertion of Peripheral Venous Catheters or Peripheral Arterial Lines

- Perform [hand hygiene](#), prepare the skin with an antiseptic and wear clean disposable gloves.

7.9. Storage, Assembly or Handling Components of Intravenous (IV) Delivery Systems

- Perform [hand hygiene](#) prior to accessing IV supplies and solutions. [13.3](#)

8. PERSONAL PROTECTIVE EQUIPMENT

Personal protective equipment IS NOT the first/only strategy used to prevent the transmission of microorganisms.

Focusing only on availability and use of various protective equipment will result in less than ideal protection of all persons, including those receiving care, and staff.

Personal protective equipment (PPE) provides a physical barrier between the uninfected and an infectious agent/infected source. It protects the user from exposure to bloodborne and other microorganisms (germs) (e.g., sprays of blood, body fluids, respiratory tract or other secretions or excretions).

Appropriate PPE must be available for use to prevent exposure to an infectious agent/infected source. Effective and appropriate use of PPE is reliant on the user's adherence and competence. Health care workers HCWs should determine what PPE is needed by performing a [Point of Care Risk Assessment \(PCRA\)](#).

Appropriate and proper use of PPE includes:

- PCRA to determine need for PPE
- Correct technique for donning and doffing PPE
- Correct technique when wearing PPE (e.g., not contaminating self)
- Discard into designated receptacles immediately after use, followed by [hand hygiene](#), preferably with ABHR

Following the [PCRA](#), required PPE may include:

- Gloves
- Gowns
- Facial protection
 - Masks (medical)
 - Eye protection (safety glasses, goggles or face shields)

NOTE: prescription or fashion glasses are NOT considered eye protection
 - Masks with visor attachment.

Over-reliance on PPE may result in a false sense of security; misuse (e.g., surgical caps and bouffants when caring for a patient/resident/ client [P/R/C] with lice); or increased waste. Putting on or removing PPE incorrectly can result in inadvertent exposure of the user or the P/R/C to infectious agents or contamination of the healthcare (external) environment. Faith or cultural head coverings shall be covered in areas where hair must be covered (e.g., Operating Room bouffant cap) but do not require covering or removing in isolation rooms.

8.1. Gloves

The use of gloves is NOT a substitute for hand hygiene, but an additional measure of protection.

For Routine Practices, glove use is dependent on a [PCRA](#) of the P/R/C, the environment and the interaction. Gloves are not required for routine care activities when contact is limited to the intact skin of the P/R/C.

Gloves include:

- Procedure
- Surgical (i.e., Sterile).

Gloves are used to reduce the transmission of microorganisms from one person to another or from one body site to another, and to reduce the risk of exposure of staff to blood, body fluids, secretions and excretions, mucous membranes, draining wounds or non-intact skin and for handling items or touching surfaces visibly or potentially soiled. **Hand hygiene is ALWAYS necessary after the removal of gloves**, as they may have microscopic holes, or hands may become contaminated during glove removal.

Wear gloves as determined by the [PCRA](#):

- For anticipated contact with blood, body fluids, secretions and excretions, mucous membranes, draining wounds or non-intact skin (including skin lesions or rash)
- For handling items or touching surfaces visibly or potentially soiled with blood, body fluids, secretions or excretions
- While providing direct care if the user has an open cut or abrasions on the hands. If gloves are used for this reason they should be changed every time hand hygiene is required.

8.1.1. Appropriate Glove Use

- Perform [hand hygiene](#) prior to putting on gloves for tasks requiring clean, aseptic or sterile technique
- Put gloves on directly before contact with the P/R/C or just before the tasks or procedure requiring gloves

- Ensure gloves are the correct size to maximize protection, dexterity and comfort. [13.3](#)
- Select type of glove appropriate to the task. [13.1](#)
- Wear disposable procedure or surgical gloves or reusable utility gloves for cleaning the environment or medical equipment. [13.1](#) If using reusable utility gloves for cleaning of the environment or medical equipment be sure to disinfect with a healthcare approved disinfectant after the task and allow to air dry away from sources of contamination
- **DO NOT reuse single use gloves. DO NOT clean gloves with alcohol-based hand rub or wash for reuse.** Washing affects integrity and has not been shown to be effective in removing microorganisms
- Remove gloves and perform [hand hygiene](#) immediately after care activities. If gloves are still indicated, replace with a clean pair
- Remove gloves and dispose into a hands-free waste receptacle immediately following their intended use. Follow immediately with [hand hygiene](#).
- Change gloves between the care of each P/R/C.

DO NOT DOUBLE GLOVE

Wearing extra PPE may affect fit and complicates the doffing process which may increase the risk of self-contamination.

To reduce hand irritation related to gloves:

- Wear gloves for as short a time as possible
- Ensure hands are clean and dry before putting on gloves
- Ensure gloves are intact and clean and dry inside.

8.2. Long-Sleeved Gowns and Other Apparel

Long sleeved cuffed gowns are worn for Routine Practices as indicated by the [PCRA](#):

- During procedures and patient/resident/client care activities likely to soil clothing and/or generate splashes or sprays of blood, body fluids, secretions or excretions
- To protect uncovered skin
- To prevent soiling of clothing.

Gowns include:

- Isolation gown
- Reusable/disposable
- Fluid repellent/resistant
- Sterile.

The type of gown selected is based on the:

- Anticipated degree of contact with infectious material
- Potential for blood and body fluid penetration of the gown (fluid repellence/resistance when heavy liquid contamination is anticipated (e.g., operating theatre, dialysis)
- Requirement for sterility (e.g., operating theatre, central line insertion).

8.2.1. Appropriate Gown Use

- Perform [hand hygiene](#) before putting on a gown
- Ensure gown is long enough to cover the front and back of the user, from the neck to mid-thigh and the sleeves no shorter than just above the wrist
- Put gown on with the opening at the back, with edges overlapping, thus covering as much clothing as possible
- Ensure cuffs of the gown are covered by gloves
- Tie the gown at the neck and waist
- Remove gown by undoing the neck and waist ties, starting with neck ties, and remove the gown without touching the clothing or agitating the gown unnecessarily; then turn the gown inside on itself, and roll it up
- Remove gown immediately after the indication for use and place in a hands-free waste receptacle (if disposable), or in a soiled linen bag (if reusable), and perform hand hygiene before leaving the care environment
- Remove wet gowns immediately to prevent a wicking action that facilitates the passage of microorganisms through the fabric
- **DO NOT** reuse gowns once removed, even for repeated contacts with same P/R/C
- **DO NOT** wear the same gown between successive P/R/Cs
- Perform [hand hygiene](#) after removing the gown due to possible contamination of hands during removal of the gown.^{[13.3](#)}

There is no evidence the routine use of gowns for all P/R/C care is beneficial in the prevention of HAIs, even in high risk units such as intensive care or haematopoietic stem cell transplant units.

Universal gown use has had no effect on HAI rates in neonatal or paediatric ICUs or on rates of neonatal colonization on post-partum wards.

In the laboratory setting, wearing of laboratory coats is considered PPE. PPE worn inside the laboratory setting should not be worn outside the laboratory containment area (e.g., should not be worn in cafeteria, lunchroom, or P/R/C areas). [13.10](#)

Several gown sizes should be available in a health care setting to ensure appropriate coverage of staff. [13.8](#)

Staff Apparel / Uniform Considerations

For aesthetic purposes and professional etiquette, staff apparel and uniforms shall be clean.

- Sleeves shall not interfere with or become wet when performing hand hygiene
- It is safe to launder staff uniforms at home
- Adhere to organizational policies regarding the laundering of scrub suits and uniforms supplied by the organization
- Personal clothing that cannot be completely covered by surgical attire shall not be worn by staff required to perform a surgical scrub.

Outside of the laboratory setting, apparel such as uniforms, laboratory coats or scrub suits may be worn by staff for purposes of comfort, convenience or identity, but DO NOT have a role in prevention of infection (i.e., they are not considered PPE).

8.3. Facial Protection

Facial protection includes medical masks (procedure or surgical mask), eye protection (safety glasses, goggles or face shields).

8.3.1.(Medical) Masks

Medical masks include procedure or surgical masks, and have several uses:

- To protect from sprays or splashes
- As a barrier for and from infectious sources
- As a barrier when performing aseptic/sterile procedure.
- To protect susceptible hosts when within two metres/six feet of patient/resident /client (P/R/C) with respiratory signs/symptoms.

8.3.2. Eye Protection

The eye is an important portal of entry for some pathogens. Pathogens may be introduced into the eye directly via respiratory droplets generated during coughing or suctioning, or by self-inoculation if the eyes are touched with contaminated fingers.

Eyes may be protected through use of regionally approved:

- Safety glasses,
- Goggles or
- Face shields.

Users should avoid touching their faces with their hands during P/R/C care. [13.3](#)

The need for facial protection during routine care is determined by the [PCRA](#):

- Interactions involving activities likely to generate coughing, splashes or sprays of blood, body fluids, secretions or excretions
- Procedures that potentially expose the mucous membranes of the eyes, nose or mouth, require facial protection. Transmission of hepatitis C and HIV has been reported by splashes of blood to the mucous membranes of the face
- When caring for a coughing and sneezing P/R/C

8.3.2.1. Appropriate Use of Facial Protection

- Perform hand hygiene before putting on facial protection
- Put on facial protection immediately before the activity that requires you to wear a medical mask and eye protection
- Remove medical mask and eye protection immediately after the activity for which it is used
- Perform [hand hygiene](#) prior to putting on facial protection
- Users should avoid touching their faces with their hands during P/R/C care [13.3](#)
- Wear and discard facial protection appropriately to prevent self-contamination
- Ensure nose, mouth and chin are covered when wearing a medical mask
- Avoid self-contamination by not touching facial protection on its external surface during use and disposal
- Wear disposable eye protection or face shields only once to avoid self-contamination

- When eye protection is required, wear eye protection over prescription or fashion glasses; ***prescription or fashion glasses alone are NOT adequate for eye protection***
- Remove facial protection carefully by the straps or ties. Bend forward to allow the medical mask to fall away from the face [13.1](#)
- Discard facial protection immediately after the intended use into a hands-free waste receptacle (i.e., dispose of as soon as removed from the face) and perform hand hygiene
- If eye protection or face shields are reusable, clean and disinfect as per organizational policy before reuse
- ***DO NOT dangle a medical mask around the neck or ears when not in use***
- ***DO NOT reuse medical mask***
- ***DO NOT place on top of head or around the neck for later use***
- Change the medical mask if it becomes wet or soiled (from the wearer's breathing or due to an external splash)
- Change the medical mask if breathing becomes difficult
- ***Do not*** fold or store medical mask in a pocket. [13.3](#)

8.4. Respiratory Protection

Respiratory protection from airborne infection requires the use of a respirator with NIOSH-approved N95 or higher filtration to prevent inhalation of microorganisms. Respiratory protection may be necessary as a component of airborne precautions or recommendations for performing [AGMPs](#) on certain P/R/C. The need for respiratory protection is determined by a [POINT OF CARE RISK ASSESSMENT \(PCRA\)](#). Factors to be considered are the specific infectious agent, known or suspected, infection status of the P/R/C, the care activity to be performed, the immune status of staff involved in care and the ability of the P/R/C to perform respiratory hygiene. [13.1](#)

9. SPECIMEN COLLECTION

All clinical specimens are considered potentially infectious and shall be handled carefully to prevent contamination.

All specimens submitted to the laboratory for testing must be packaged in such a manner as to prevent spillage, breakage, or damage to the specimen itself, and/or to accompanying specimens. The safety of the environment, and the safety of all persons involved in the shipping, handling and receiving of these specimens must be ensured by preventing exposure to the contents of the shipment at any time.

Specimens and requisitions must be labelled to comply with receiving laboratory's acceptance policy.

Specimens shall be transported to the laboratory in sealable zipper storage bags (e.g., Ziploc®).

Each specimen must have its own requisition. Requisitions shall be placed in the exterior pouch of the sealable zipper storage bag (e.g., Ziploc®) for transport.

Consider [Personal Protective Equipment](#) when collecting specimens.

Perform [hand hygiene](#) immediately before and after specimen collection.



9.1. Process

Specimens will be rejected for analysis for the following reasons:

- Specimens that cannot be safely processed, i.e., specimens with needle attached
- Improperly transported specimens
- Improperly labelled specimens
- Improper specimen collection
- Samples that are inappropriate for the test requested

**Specific rejection criteria exist depending on the specimen type;
 Consult the laboratory for further information**

10. SHARPS SAFETY & PREVENTION OF BLOODBORNE TRANSMISSION

Adherence to Routine Practices can reduce the transmission of microorganisms (germs). As such, Additional Precautions are not routinely required for the care of people with bloodborne pathogens such as HIV and hepatitis

The prevention of sharps injury and staff exposure to bloodborne pathogens is a component of Routine Practices.

Users of sharps require education and training about how to safely handle sharp devices to prevent injuries to themselves and to others who may encounter the device during or after procedures. Safety programs include a formal incident investigation for every sharp injury occurring in the work setting.

Use of safety engineered devices such as using protected needle devices, needle-free systems with self-sealing ports, and syringes with safety features, have been reported to reduce needlestick injuries. Their use has been identified as a priority in risk reduction strategies. Some models have demonstrated a risk for patient/resident/client (P/R/C). Therefore, careful consideration to both the P/R/C and staff should be taken when selecting safety engineered sharps devices. Refer to [WRHA policy 20.20.020, Sharps, Safe Handling, Use and Disposal \(including Safety-Engineered Needle \(SEN\) Exemption\)](#).

DO NOT recap used needles. Handle used needles and other sharp instruments with care to avoid injuries during disposal. Dispose of used needles and other used single use sharp items immediately into designated puncture-resistant containers readily accessible at the point of care.

In home care settings:

- Teach P/R/C and caregivers in the home the correct procedures for safe handling and disposal of sharps and sharp containers in according to municipal or regional regulations
- Insure home storage of sharps is in a labelled, puncture-proof container with a tight-fitting lid that prevents leakage
- Sharps containers may be free at some pharmacies or can be for purchase from many pharmacies or medical supply stores.

Protect eyes, nose and mouth (using facial protection) when splashes with blood and/or body fluids are anticipated.

Perform first aid **immediately** if exposed to blood or body fluids:

- Thoroughly rinse the site of a percutaneous injury with running water and gently clean any wound with soap and water
- Flush mucous membranes of the eyes, nose, or mouth with running water if contaminated with blood, body fluids, secretions or excretions
- Thoroughly rinse non-intact skin with running water if contaminated with blood, body fluids, secretions or excretions.

Report immediately to employer after first aid and seek immediate medical attention. **Refer to [WRHA Occupational and Environmental Safety and Health Operational Procedure, Blood and Body Fluid – Post Exposure Management](#).**

11. MANAGEMENT OF THE PATIENT/RESIDENT/CLIENT CARE ENVIRONMENT

A clean environment is a safer environment for patients/residents/clients (P/R/C) in all health care settings. The risk of healthcare-associated infections (HAIs) can be reduced when surfaces, items, and equipment are cleaned and disinfected with the correct products and at the right times.

11.1. Cleaning of Environment

Minimize Environmental Contamination:

- **DO NOT** bring the care record/chart into the P/R/C room, cubicle or designated bed space in a shared room and perform hand hygiene after handling the record/chart
- **DO NOT** eat or drink in areas where direct care is provided, at the nursing station, in medication rooms, in clean supply rooms, and in reprocessing or laboratory areas
- Dedicate non-critical medical equipment to a single P/R/C
- Assign responsibility and accountability for routine cleaning and disinfection of care equipment
- Ensure environmental cleaning follows a set procedure and frequency, and is documented and supervised by adequately trained dedicated personnel
- Ensure adequate human resources^{[13.8](#)}

Areas/programs should have:

- ☒ Written policies and procedures for cleaning and disinfection of patient/resident/client rooms and equipment that includes cleaning standards and frequencies^{[13.8](#)}
- ☒ Procedures and increased capacity for outbreak management^{[13.8](#)}
- Ensure surfaces are constructed of materials that can be easily cleaned at the point of use

- Increase the frequency of cleaning and disinfecting high touch surfaces. Clean and disinfect surfaces likely to be touched and/or used on a more frequent schedule compared to other surfaces (high touch surfaces).
 - This includes surfaces in close proximity to the P/R/C (e.g., bedrails, over bed tables, call bells, exam beds, treatment chairs) and frequently touched surfaces in the care environment such as door knobs, surfaces in the P/R/C's bathroom and shared common areas for dining, bathing, toileting
 - This also includes personal use items such as stethoscopes, wipeable lanyards and pens.
- Monitor for adherence to recommended environmental cleaning practices
- Ensure rooms/spaces are terminally cleaned following P/R/C appointment, discharge and after discontinuing precautions
- Use facility approved cleaners and disinfectants
- Ensure the availability of healthcare approved cleaners and disinfectants for Environmental Services Staff and frontline staff [13.4](#)
- Clean areas adjacent to construction activities at the end of the day or at other times as required in order to maintain cleanliness of the area. [13.8](#)

In outbreak situations, or when continued transmission of certain microorganisms (e.g., norovirus, rotavirus, *C. difficile*) occurs, specific disinfectant products may need to be used as appropriate (i.e., facility/regionally approved sporicidal agents). Regional IP&C approval is required prior to use of these specialized products.

11.2. Cleaning and Disinfection of Non-Critical Patient/Resident/Client Care Equipment

Contamination of care equipment and items in the care environment, as well as the care environment itself has been implicated in infection transmission. Follow policies and procedures for containing, transporting, and handling used patient/resident/client (P/R/C) care equipment and medical instruments and devices

Clean and disinfect all non-critical re-useable equipment when soiled and between uses with different P/R/Cs. [13.8](#)

Identify used non-critical care equipment and other items such as toys and electronic games, and **DO NOT** allow use by another P/R/C until these items are appropriately cleaned and disinfected.

Clean and disinfect non-critical care equipment dedicated to an individual P/R/C when soiled and on a regular schedule.

Dedicate bedpans and commodes to each P/R/C and label appropriately. Clean and disinfect before use by another P/R/C. The use of disposable bedpans is acceptable. Bedpan holders for disposable bedpans must be reprocessed following use.

Store sterile and clean supplies in a designated and separate clean dry area protected from dust. **DO NOT** store under sinks and/or near plumbing as leaks may occur.

Discard personal care items (e.g., tissues, lotions, soaps, razors) and disposable equipment such as containers used for blood collection or tourniquets left in the room following transfer, terminal cleaning or discharge.

Unless a computer keyboard and computer device technology has been just cleaned/disinfected by the user, consider keyboards and devices used in the healthcare (external) environment as contaminated. Clean hands after using keyboards and computer devices. Assign responsibility for regular cleaning and disinfection of computer keyboards and horizontal computer cart surfaces utilized in the healthcare (external) environment.

Ensure computer keyboards in P/R/C rooms are cleaned and disinfected after each use, and upon discharge or during terminal cleaning/disinfection.

11.2.1. In Home Care Settings

Educate clients about the importance of environmental cleaning.

Limit the amount of disposable and non-disposable care equipment and supplies brought into the home.

Advise clients to purchase items such as thermometers and scissors for personal use whenever possible.

Leave reusable care equipment in the home until the person is discharged from home care services whenever possible.

Clean and disinfect non-critical care equipment (e.g., stethoscope) that cannot remain in the home before removal from the home.

Alternatively, place contaminated reusable items in a plastic bag for transport then cleaned and disinfected in a designated area at the home care office.

Where limiting supply entering home is not possible, prior to entry, supplies which are not in sealed impervious packaging can be bagged and sealed to prevent contamination and support retrieval and reprocessing if appropriate and needed

Avoid opening new packaging to decant into client's home environment, only open as needed prior to use

Removal for reprocessing of supplies shall not occur in the following instances:

- a. Client(s) home has a known or suspected bed bugs, rodent activity
- b. Client or housemate has a communicable disease
- c. Client or housemate has an infectious process requiring Additional Precautions,
- d. Hoarding environment present
- e. Presence of gross environmental soiling with blood or body fluids
- f. Supplies unable to withstand cleaning and disinfection:
 - Paper packaging
 - Opened supplies
- g. Expired supplies.

11.3. Handling of Linen

Linen in healthcare facilities may become contaminated with pathogens; risk of disease is usually negligible

Care should be taken in the handling of soiled linen to prevent dispersal of microorganisms. Handle soiled linen with a minimum of agitation to avoid contamination of air, surfaces, self and other persons.

Place soiled linen in a no-touch receptacle at the point of use.

Use leak-proof containers for laundry contaminated with blood or bodily secretions (urine, feces, etc.). Water soluble bags and 'double-bagging' are not recommended. [13.8](#)

Tie linen bags securely and **DO NOT** over-fill [13.8](#)

Transported and store clean linen in a manner to prevent inadvertent handling or contamination by dust, which may contain fungal spores harmful to immunocompromised P/R/Cs.

Maintain separation of clean and soiled linen during transport and storage.

If laundry chutes are used, they should be properly designed, maintained, and used in a manner to minimize dispersion of aerosols from contaminated laundry.

Change linen regularly and when soiled, upon discontinuation of Contact Precautions and following discharge/transfer of the P/R/C. In ambulatory care/clinic areas change linen following every treatment/procedure.

Roll or fold heavily soiled linen to contain the heaviest soil in the centre of the bundle. **DO NOT** spray soiled linens with water; use a gloved hand and toilet tissue to remove any solid waste. To avoid splashing carefully place into a bedpan or toilet for flushing.

Perform hand hygiene after handling soiled linen.

Wash reusable linen bags after each use; they may be washed in the same cycle as the linen contained in them.

11.4. Handling of Waste

Most waste generated in healthcare settings is no more hazardous than household waste.

Waste receptacles should be conveniently located and, preferably, hands-free.

DO NOT double-bag waste unless the first bag becomes stretched or damaged, or when waste has spilled on the exterior. [13.3](#)

Close waste bags when three-quarters full and tie in a manner that prevents contents from escaping. [13.3](#)

Remove waste to central holding areas at frequent intervals. [13.3](#)

Dispose of blood, suctioned fluids, excretions and secretions in a sanitary sewer or septic system according to municipal/regional regulations. [13.1](#)

Contain and dispose of biomedical, pharmaceutical and sharps waste according to site policies.

Wear personal protective equipment according to PCRA.

Perform hand hygiene after handling waste and waste containers.

11.5. Handling of Dishes

There are no indications for the use of disposable dishes other than when dishwashing equipment is non-functioning.

Perform hand hygiene after handling dirty dishes.

11.6. Handling of Deceased Bodies

Use Routine Practices properly and consistently for the routine handling of deceased bodies. There are no special requirements when handling deceased bodies. Adhere to provincial specified communicable disease regulations, available at Province of Manitoba, [Public Health Act, Dead Bodies Regulation](#).

12. VISITOR AND DESIGNATED CAREGIVER MANAGEMENT AND EDUCATION

Visitors have been documented to transmit infections including tuberculosis, pertussis, and respiratory viruses in healthcare settings.

***Visitors** - Family and friends who visit for social reasons. Their time with the patient/resident/client is discretionary and short term. They are not involved in the care of the patient/resident/client.

***Designated Caregivers (DC)** - provide physical, psychological and emotional support, as deemed important by the P/R/C. This care can include support in decision making, care coordination and continuity of care. Designated Caregiver can include family members, close friends or other caregivers and are identified by the P/R/C or substitute decision maker.

NOTE: Designated Caregiver language replaces essential care partner and designated family caregiver.

Visitors and Designated Caregivers have a responsibility to comply with Routine Practices. All staff involved in care is responsible to teach those receiving care and visitors basic principles, such as [hand hygiene](#), [respiratory hygiene](#), and use of [personal protective equipment](#).

Visiting policies must balance the risk of transmission of infectious diseases and the promotion of patient/resident/client and family centered care. Exclusion of those with signs and symptoms of transmissible infections should reduce this risk. For essential visits (e.g., parent, guardian or designated caregiver), instruct the visitor with an infection/signs and symptoms of an acute infection (e.g., cough, fever, vomiting, diarrhea, coryza, rash, conjunctivitis) on measures to take to reduce the risk of transmission (e.g., wear a medical mask for a respiratory tract infection, perform appropriate hand hygiene, remain in the P/R/C's room, avoid public areas, avoid contact with other P/R/Cs or with care equipment).

Visitors and Designated Caregivers could be at risk for serious diseases should they acquire the infection of the P/R/C (e.g., acquisition of a respiratory virus by a visitor with chronic lung disease, or exposure of a non-immune visitor to varicella). They should be capable of complying with the necessary precautions to prevent indirect transmission to others receiving care (e.g., hand hygiene, not sharing personal items).

Provide education to P/R/Cs, their families and visitors regarding [respiratory hygiene](#) and [hand hygiene](#).

13. REFERENCES

- 13.1 [Routine Practices and Additional Precautions: Preventing the Transmission of Infection in Healthcare](#) (2019, June). Manitoba Health. Accessed July 6, 2022
- 13.2 [Hand Hygiene in Outpatient and Home-Based Care and Long-Term Care Facilities](#) (2012). World Health Organization (WHO). Accessed October 26, 2022
- 13.3 [Infection Prevention and Control for Clinical Office Practice](#) (April 2015). Public Health Ontario (PHO). Provincial Infectious Disease Advisory Committee (PIDAC). Accessed July 6, 2022
- 13.4 [Best Practices for Hand Hygiene in All Health Care Settings, 4th ed.](#) (April 2014). Public Health Ontario (PHO). (PIDAC). Accessed October 26, 2022
- 13.5 [IPAC Canada Practice Recommendations - Hand Hygiene in Health Care Settings](#) (October 2022). Infection Prevention & Control Canada (IPAC Canada). Accessed December 22, 2022
- 13.6 [Best Practices for Hand Hygiene in All Health Care Settings, 4th ed.](#), (April 2014). PIDAC Accessed January 15, 2023.
- 13.7 [Provincial Guidance for Aerosol Generating Medical Procedures \(AMGPs\)](#) August 19, 2022. Shared Health Manitoba.
- 13.8 [Routine Practices and Additional Precautions in All Health Care Settings, 3rd Edition](#) (2012, November). Provincial Infectious Disease Committee (PIDAC).
- 13.9 [Hand Hygiene Practices in Healthcare Settings](#) (2012). Public Health Agency of Canada (PHAC). Accessed July 6, 2022
- 13.10 [Routine Practices and Additional Precautions for the Preventing the Transmission of Infection in Healthcare Setting](#) (2017, August). Public Health Agency of Canada (PHAC). Accessed July 4, 2023
- 13.11 Infection Prevention & Control Program Team Expert Opinion (May 2021)
- 13.12 [Purell Advanced Hand Rub Brochure](#). (2020) Gojo industries Inc.

Appendix A Surgical Hand Asepsis

The goal of surgical hand asepsis is to remove the number of transient flora and reduce resident flora. It shall be done prior to participating in a surgical procedure.⁽¹⁾ Even after skin asepsis skin is not considered sterile, rather it is deemed surgically clean.

Surgical hand asepsis is the process of removing microorganisms from the hands and forearms using either mechanical washing with a surgical scrub agent (surgical hand scrub) or chemical antisepsis with an alcohol based hand rub (surgical hand rub).⁽¹⁾ Due to the superior antimicrobial activity ABHR (surgical hand rub) is the preferred method of preoperative surgical hand preparation.⁽²⁾ Many formulations also contain long acting compounds such as chlorhexidine gluconate.⁽²⁾

Surgical hand asepsis is only effective if the hands are free from hand and arm jewelry as well as watches so the product can reach all surfaces of the hands and forearms.

Surgical hand asepsis procedure (using either a scrub or rub) should follow a standardized protocol established and approved by the healthcare organization and follow the manufacturer's instructions for use.⁽³⁾

Any product used for surgical hand asepsis should:⁽²⁾

- ☒ Inhibit the growth of microorganisms under gloved hands
- ☒ Have as wide a spectrum for antimicrobial activity as possible
- ☒ Have a prolonged antiseptic effect
- ☒ Have a persistent antimicrobial activity due to rapid multiplication of bacteria under surgical gloves and the chance of glove punctures during surgery.

REFERENCES:

1. [Surgical Aseptic Technique and Sterile Field](#). (June 30, 2023). Alberta Health Services (AHS). Accessed August 24, 2023.
2. [Best Practices for Hand Hygiene in All Health Care Settings, 4th ed.](#) (April 2014). Public Health Ontario (PHO). (PIDAC). Accessed August 24, 2023
3. The ORNAC Standards, Guidelines, and Position Statements for Perioperative Registered Nurses. 15th edition. (April 2021). Operating Room Nurses Association of Canada (ORNAC). Accessed August 24th, 2023.